

Decay Properties of the Connectivity for Mixed Long Range Percolation Models on $\mathbb{Z}^d$

Gastão A. Braga, Leandro M. Ciolleti and Rémy Sanchis

Departamento de Matemática - UFMG

Caixa Postal 1621 30161-970 - Belo Horizonte - MG - Brazil

May 10, 2007

Abstract

In this short note we consider mixed short-long range independent bond percolation models on $\mathbb{Z}^{k+d}$. Let p_{uv} be the probability that the edge (u, v) will be open. Allowing a x, y -dependent length scale and using a multi-scale analysis due to Aizenman and Newman, we show that the long distance behavior of the connectivity τ_{xy} is governed by the probability p_{xy} . The result holds up to the critical point.

1 Introduction

In this short note we consider a long range percolation model on $\mathbb{L} = (\mathbb{Z}^{k+d}, \mathbb{B})$, where $u \in \mathbb{Z}^{k+d}$ is of the form $u = (\vec{u}_0, \vec{u}_1)$, with $\vec{u}_0 \in \mathbb{Z}^k$ and $\vec{u}_1 \in \mathbb{Z}^d$ and $\mathbb{B}$ is the set of edges (unordered pairs) (u, v) , $u \neq v \in \mathbb{Z}^{k+d}$. To each edge (u, v) we associate a Bernoulli random variable ω_b which is open ($\omega_b = 1$) with probability

$$p_{uv} = p_{uv}(\beta) \equiv \beta J_{uv}, \quad u, v \in \mathbb{Z}^{k+d} \quad (1)$$

where $\beta \in [0, 1]$ and, for $\epsilon > 0$, J_{uv} is

$$J_{uv} = \begin{cases} 2(1 + \|\vec{u}_1 - \vec{v}_1\|^{d+\epsilon})^{-1} & \text{if } \vec{u}_0 = \vec{v}_0 \text{ and } \vec{u}_1 \neq \vec{v}_1; \\ 1 & \text{if } \vec{u}_1 = \vec{v}_1 \text{ and } \|\vec{u}_0 - \vec{v}_0\| = 1; \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

We denote the event $\{\omega \in \Omega : \text{there is an open path connecting } x \text{ to } y\}$ by $\{x \leftrightarrow y\}$ and define the connectivity function by $\tau_{xy} \equiv P\{x \leftrightarrow y\}$. Let $\|x\| = |x_1| + \dots + |x_d|$ be the L^1 norm on $\mathbb{Z}^d$ and $\beta_c = \sup\{\beta \in [0, 1] : \chi(\beta) < \infty\}$. Our aim is to show the following result

Theorem. *Suppose $\beta < \beta_c$ and consider the long range percolation model with p_{uv} given by (1) and J_{uv} given by (2). Then there exist positive constants $C = C(\beta)$ and $m = m(\beta)$ such that*

$$\tau_{xy} \leq \frac{C e^{-m\|\vec{x}_0 - \vec{y}_0\|}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} \quad (3)$$

for all $x, y \in \mathbb{Z}^{k+d}$.

The above result says that the probability p_{uv} dictates the long distance behavior of the connectivity function in the subcritical regime (similar lower bounds are easily obtained from the FKG inequality). For the one dimensional $\mathbb{Z}^{0+1}$ percolation model, the above result is known to hold, see [1], the same being true for one dimensional $\mathbb{Z}^{0+1}$ $O(N)$ spin models, $1 \leq N \leq 4$, see [2]. The result is expected to hold in the d -dimensional $\mathbb{Z}^{0+d}$ lattice but it is not clear how to prove it if $\beta < \beta_c$, although one can see it holds if $\beta \approx 0$. Our upper bound (3) holds if $\beta < \beta_c$ and for $(k+d)$ -dimensional lattices, $k \geq 0$ and $d \geq 1$. For lattice spin models, the upper bound (3) is known at the high temperature regime, see Ref. [3] for bounded spin models and Ref. [4] for unbounded (and discrete) ones. Ref. [5] extends some of the results of [3, 4] to a general class of continuous spin systems, with J_{uv} given by (1) and $u, v \in \mathbb{Z}^{0+d}$, while [6] considers the more general mixed decay model. In both cases, the polymer expansion (see [7] and references therein) is used.

The mixed percolation model we study is motivated by equivalent ferromagnetic models that arise as special cases of discrete space-time field theories as in [6] where the discretized Laplacian is replaced by a nearest neighbor ferromagnetic interaction. The model is interesting by itself and it has already appeared in the literature [8].

The Hammersley-Simon-Lieb inequality [9, 10, 11] is a key ingredient in [1] and [2] and here we also adopt this ‘‘correlation inequality’’ point of view. For completeness, we state this inequality in the form we will use, see [12]. For each set $S \subset \mathbb{Z}^d$, let $\tau_{xy}^S \equiv P\{x \leftrightarrow y \text{ inside } S\}$. Then

Hammersley-Simon-Lieb Inequality (HSL) *Given $x, y \in \mathbb{Z}^{k+d}$, if $S \subset \mathbb{Z}^{k+d}$ is such that $x \in S$ and $y \in S^c$, then*

$$\tau_{xy} \leq \sum_{\{u \in S, v \in S^c\}} \tau_{xu}^S p_{uv} \tau_{vy}.$$

We now recall some known facts about the long range percolation model defined by (1) and (2). Let $\theta(\beta, \varepsilon) = P_{\beta, \varepsilon}\{0 \leftrightarrow \infty\}$ be the probability that the origin will be connected

to infinity. If $k + d \geq 2$ is the space dimension then, by comparing with the nearest neighbor model and for any positive ε , there exists $\beta_c = \beta_c(d, \varepsilon)$ such that $\theta(\beta, \varepsilon) = 0$ if $\beta < \beta_c$ and $\theta(\beta, \varepsilon) > 0$ if $\beta > \beta_c$. For $k = 0$ and $d = 1$, it is known that the existence of β_c depends upon ε , if $\varepsilon > 1$ then there is no phase transition [13] while it shows up if $0 \leq \varepsilon \leq 1$, see [14]. A phase transition can also be measured in terms of χ , the mean cluster size, given by $\chi = \sum_x \tau_{0x}$. Let $\pi_c(d, \varepsilon) = \sup\{\beta : \chi(\beta, \varepsilon) < \infty\}$. Then, it comes from the FKG Inequality [15] that $\pi_c(d, \varepsilon) \leq \beta_c(d, \varepsilon)$. The equality $\pi_c(d, \varepsilon) = \beta_c(d, \varepsilon)$ holds for the class of models we are dealing with and it was proved independently by Aizenman and Barsky in [16] and Menshikov in [17]. We will use the condition $\chi < \infty$ to characterize the subcritical region.

The remaining of this note is divided as follows: in the next section we prove Theorem 1 and in Section 3 we make some concluding remarks regarding the validity of our results to ferromagnetic spin models.

2 Proof of the Theorem

Let $x = (\vec{x}_0, \vec{x}_1) \in \mathbb{Z}^{k+d}$. We first observe that

$$\chi = \sum_{n \geq 0} \sum_{\|\vec{x}_0\|=n} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x}.$$

Since $\chi < \infty$, given $\lambda \in (0, 1)$, there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$, we have

$$\sum_{\|\vec{x}_0\|=n} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} < \lambda.$$

Consider now $x = (\vec{x}_0, \vec{x}_1)$ with $\|\vec{x}_0\| > n_0$. Using the translation invariance of the model and applying iteratively the HSL Inequality with $S = \{x \in \mathbb{Z}^{k+d}; \|\vec{x}_0\| \leq n_0\}$, we obtain

$$\sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} \leq \lambda^{\lfloor \|\vec{x}_0\|/n_0 \rfloor} \leq C_1 \exp(-(m + \delta)\|\vec{x}_0\|),$$

where $\lfloor r \rfloor$ denotes the integer part of r , $\delta > 0$ is given and m is defined by $e^{-(m+\delta)} = \lambda^{1/n_0}$. Next we show that a HSL type inequality holds for the modified connectivity function $T_m(x, y) \equiv e^{m\|\vec{x}_0 - \vec{y}_0\|} \tau_{xy}$ and for the set $S = \mathcal{C}_r(x) \equiv \{z \in \mathbb{Z}^{k+d}; \|\vec{x}_1 - \vec{z}_1\| \leq r\}$. Applying the HSL Inequality with the above specified S , we have

$$\tau_{xy} \leq \sum_{\substack{u \in \mathcal{C}_r(x) \\ v \in \mathcal{C}_r^c(x)}} \tau_{xu} p_{uv} \tau_{vy}.$$

Then, for $y \in \mathcal{C}_L^c(x)$ for some $L > 1$, we obtain

$$\begin{aligned} T_m(x, y) &= e^{m\|\vec{x}_0 - \vec{y}_0\|} \tau_{xy} \leq e^{m\|\vec{x}_0 - \vec{y}_0\|} \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x)}} \tau_{xu} p_{uv} \tau_{vy} \\ &\leq \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x)}} e^{m\|\vec{x}_0 - \vec{u}_0\|} \tau_{xu} p_{uv} e^{m\|\vec{y}_0 - \vec{v}_0\|} \tau_{vy} \end{aligned}$$

since we necessarily have that $\vec{u}_0 = \vec{v}_0$. It then follows that

$$T_m(x, y) \leq \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x)}} T_m(x, u) p_{uv} T_m(v, y).$$

We remark that $\chi_m \equiv \sum_{x \in \mathbb{Z}^{k+d}} T_m(0, x) < \infty$ is finite if $\beta < \beta_c$ since

$$\begin{aligned} \sum_{x \in \mathbb{Z}^{k+d}} T_m(0, x) &= \sum_{x \in \mathbb{Z}^{k+d}} e^{m\|\vec{x}_0\|} \tau_{0x} \leq \sum_{k \geq 0} \sum_{\|\vec{x}_0\|=k} e^{m\|\vec{x}_0\|} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} \\ &\leq \sum_{k \geq 0} 2dk^{d-1} e^{-\delta k} < \infty. \end{aligned}$$

From now on we closely follow Section 3 of [1] and prove the polynomial decay of T_m up to the critical point. For fixed $x, y \in \mathbb{Z}^{k+d}$ and $L \equiv \|\vec{x}_1 - \vec{y}_1\|/4$, we know that

$$\begin{aligned} T_m(x, y) &\leq \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x)}} T_m(x, u) p_{uv} T_m(v, y) \\ &\leq \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x) \cap \mathcal{C}_{3L}(x)}} T_m(u, x) p_{uv} T_m(v, y) + \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x) \cap \mathcal{C}_{3L}^c(x)}} T_m(x, u) p_{uv} T_m(v, y). \end{aligned} \quad (4)$$

Let

$$\mathbb{T}_m(L) \equiv \sup\{T_m(0, u); u \in \mathcal{C}_L^c(0)\} \quad \text{and} \quad \gamma_L \equiv \sum_{\substack{u \in \mathcal{C}_L(x) \\ v \in \mathcal{C}_L^c(x)}} T_m(x, u) p_{uv}.$$

Then, the first term on the r.h.s. of (4) is bounded above by $\mathbb{T}_m(L/2)\gamma_L$ while the second one is bounded by

$$\frac{2^{d+\varepsilon} 2\beta \chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}},$$

leading to

$$T_m(x, y) \leq \frac{2^{d+\varepsilon} 2\beta \chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \gamma_L \mathbb{T}_m\left(\frac{L}{2}\right).$$

Now, since $\chi_m < \infty$ for $\beta < \beta_c$ and since $\sum_u p_{0u} < \infty$, we have that $\gamma_L \rightarrow 0$ as $L \rightarrow \infty$. For $\alpha \in (0, 2^{-(d+\varepsilon)})$, there exists $L_0 > 0$ such that $\gamma_L < \alpha$ for all $L \geq L_0$. Considering $L > L_0$, it follows that

$$\mathbb{T}_m(L) \leq \frac{2^{d+\varepsilon} 2\beta\chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \alpha \mathbb{T}_m\left(\frac{L}{2}\right). \quad (5)$$

Iterating (5) n times, with n the smallest integer for which $L2^{-n} \leq L_0$, we have for all $L > L_0$

$$\mathbb{T}_m(L) \leq \frac{2\beta\chi_m^2 \sum_{j=0}^{n-1} (\alpha 2^{d+\varepsilon})^j}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \alpha^n \mathbb{T}_m\left(\frac{L}{2^n}\right).$$

Noting that $T_m(x, y) \leq \mathbb{T}_m(L)$, that $\mathbb{T}_m(L) \leq 1$ for any $L > 0$ and that

$$\alpha^n \leq 2^{-(d+\varepsilon)n} = \frac{1}{(1 + L^{d+\varepsilon})} \frac{(1 + L^{d+\varepsilon})}{2^{(d+\varepsilon)n}} \leq \frac{2 \cdot 2^{d+\varepsilon}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} \left(\frac{L}{2^n}\right)^{d+\varepsilon} \leq \frac{2(2L_0)^{d+\varepsilon}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}},$$

we can conclude that, for $\beta < \beta_c$,

$$\tau_{xy} \leq \frac{C e^{-m\|\vec{x}_0 - \vec{y}_0\|}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}}$$

and the bound (3) holds. □

3 Concluding Remarks

The strategy used to prove Theorem 1 can also be applied to $\mathbb{Z}^d$ ferromagnetic models with free boundary conditions and pair interaction J_{uv} given by (2). For this class of models the Griffiths inequalities [18, 19] are valid, guaranteeing the positivity of spin-spin correlations $\langle \sigma_x \sigma_y \rangle$. Simon-Lieb Inequality also holds in this case (see [7] and references therein), with p_{uv} replaced by βJ_{uv} , where β is the inverse of the temperature, and with τ_{xy} replaced by $\langle \sigma_x \sigma_y \rangle$. Finally, since the uniqueness of the critical point is guaranteed in [20], the results of Section 2 are also valid for these models.

Acknowledgements: G. B. was partially supported by CNPq; L. C. acknowledges CNPq for a graduate scholarship; R. S. was partially supported by Pró-Reitoria de Pesquisa - UFMG under grant 10023.

References

- [1] M. Aizenman and C. M. Newman, *Discontinuity of the percolation density in one-dimensional $1/|x - y|^2$ percolation models*, Comm. Math. Phys. **107** (1986), no. 4, 611–647

- [2] H. Spohn and W. Zwerger, *Decay of the two-point function in one-dimensional $O(N)$ spin models with long-range interactions*, J. Stat. Phys. **94**, 1037 - 1043 (1999).
- [3] L. Gross, *Decay of Correlations in Classical Lattice Models at High Temperature*, Commun. Math. Phys. **68**, 9-27 , (1979).
- [4] R. Israel and C. Nappi, *Exponential Clustering for Long-Range Integer-Spin Systems*, Commun. Math. Phys. **68**, 29-37 , (1979).
- [5] A. Procacci and B. Scoppola, *On Decay of Correlations for Unbounded Spin Systems with Arbitrary Boundary Conditions*, J. Stat. Phys. **105**, 453-482 , (2001).
- [6] R. S. Thebaldi, E. Pereira and A. Procacci, *A cluster expansion for the decay of correlations of light-mass quantum crystals and some stochastic models under intense noise*, J. Math. Phys. **46**, 053303 , (2005).
- [7] J. Glimm and A. Jaffe, *Quantum Physics: A Functional Integral Point of View*, Second Edition, Springer-Verlag, New York, (1987).
- [8] V. Sidoravicius, D. Surgailis and M.E. Vares, *On the truncated anisotropic long-range percolation on Z^2* , Stochastic Processes and their Applications, **88**, 337-349, (1999).
- [9] J.M. Hammersley, *Percolation Processes. Lower bounds for the critical probability*, Annals of Mathematical Statistics. **28**, 790-795 , (1957)
- [10] B. Simon, *Correlation Inequalities and the Decay of Correlations in Ferromagnets*, Commun. Math. Phys., **77**, 111-126, (1980).
- [11] E. H. Lieb, *A Refinement of Simon's Correlation Inequality*, Commun. Math. Phys., **77**, 127-135, (1980).
- [12] M. Aizenman and C. M. Newman, *Tree graph inequalities and critical behaviour in percolation models*, J. Stat. Phys. **36**, 107-143 (1984).
- [13] L.S. Schulman, *Long range percolation in one dimension*, J. Phys. Lett. **A16** L639 (1983).
- [14] C.M. Newman, L.S. Schulman, *One-dimensional $1/|j-i|^s$ percolation models: The existence of a transition for $s \leq 2$* , Comm. Math. Phys. **104** (1986), 547-571.
- [15] G. Grimmett, *Percolation*, Second Edition, Springer-Verlag, Berlin, (1999).
- [16] M. Aizenman and D.J. Barsky, *Sharpness of the phase transition in percolation models*, Comm. Math. Phys. **108** (1987), no. 3, 489-526.
- [17] M. V. Menshikov, *Coincidence of Critical Points in Percolation Problems*, Soviet Mathematics Doklady **33** (1986), 856-859.

- [18] R. B. Griffiths, *Correlations in Ising Ferromagnets I*, J. Math. Phys., **8**, 478-483, (1967).
- [19] R. B. Griffiths, *Correlations in Ising Ferromagnets. II. External Magnetic Fields*, J. Math. Phys., **8**, 484-489, (1967).
- [20] M. Aizenman, D. J. Barsky and R. Fernández, *The phase transition in a general class of Ising-type models is sharp*, J. Statist. Phys. **47** (1987), no. 3-4, 343-374.